

PCR based mutation screening of the laminin $\alpha 2$ chain gene (LAMA2): application to prenatal diagnosis and search for founder effects in congenital muscular dystrophy

Pascale Guicheney, Nicolas Vignier, Xu Zhang, Yi He, Corinne Cruaud, Véronique Frey, Anne Helbling-Leclerc, Pascale Richard, Brigitte Estournet, Luciano Merlini, Haluk Topaloglu, Marina Mora, Jean-Paul Harpey, Charles-Antoine Haenggeli, Annie Barois, Bernard Hainque, Ketty Schwartz, Fernando M S Tomé, Michel Fardeau, Karl Tryggvason

INSERM U153,
Groupe Hospitalier
Pitié-Salpêtrière,
Institut de Myologie,
47 Boulevard de
l'Hôpital, 75651 Paris
Cedex 13, France
P Guicheney
N Vignier
Y He
A Helbling-Leclerc
K Schwartz
F M S Tomé
M Fardeau

Department of
Medical Biochemistry
and Biophysics,
Karolinska Institute,
171 77 Stockholm,
Sweden
X Zhang
K Tryggvason

Généthon, 1 Rue de
l'Internationale, 91002
Evry, France
C Cruaud

Service de Biochimie,
Groupe Hospitalier
Pitié-Salpêtrière, 47
Boulevard de
l'Hôpital, 75651 Paris
Cedex 13, France
V Frey
P Richard
B Hainque

Service de
Pédiatrie-Réanimation
Infantile, Hôpital
Raymond-Poincaré,
92380 Garches, France
B Estournet
A Barois

Neuromuscular
Laboratory, Rizzoli
Orthopaedic Institute,
Bologna, Italy
L Merlini

Department of
Paediatric Neurology,
Hacettepe Children's
Hospital, Ankara,
Turkey
H Topaloglu

Abstract

Classical congenital muscular dystrophy with merosin deficiency is caused by mutations in the laminin $\alpha 2$ chain gene (LAMA2). Extended sequencing of the introns flanking the 64 LAMA2 exons was carried out and, based on these sequences, oligonucleotide primers were designed to amplify the coding region of each exon separately. By PCR-SSCP analysis, we identified eight new mutations in nine families originating from various countries. All induced a premature truncation of the protein, either in the short arm or in the globular C-terminal domain. A 2 bp deletion in exon 13, 2098delAG, was found in three French non-consanguineous families and a nonsense mutation of exon 20, Cys³⁶stop, in two other non-consanguineous families originating from Italy. Determination of rare intragenic polymorphisms permitted us to show evidence of founder effects for these two mutations suggesting a remote degree of consanguinity between the families. Other, more frequent polymorphisms, G to A 1905 (exon 12), A to G 2848 (exon 19), A to G 5551 (exon 37), and G to A 6286 (exon 42), were used as intragenic markers for prenatal diagnosis. This study provides valuable methods for determining the molecular defects in LAMA2 causing merosin deficient congenital muscular dystrophy.

(*J Med Genet* 1998;35:211-217)

Keywords: congenital muscular dystrophy; merosin; laminin-2

Merosin deficient congenital muscular dystrophy (MD CMD) is an autosomal recessive muscle disorder with homogeneous clinical features represented by early onset of muscle hypotonia, weakness, severe contractures, and white matter changes in the central nervous system.¹⁻³ The children generally have normal intelligence and in a number of cases epileptic seizures.⁴ Muscle biopsies show marked variation in fibre size, necrotic and regenerating processes at early stages, and interstitial fibrosis. Immunohistochemical analyses have

shown absence or reduction of the laminin $\alpha 2$ chain around the skeletal muscle fibres.⁵ Disruption of the basement membrane of muscle fibres based on electron microscopy examination has been reported in a few cases.⁶⁻⁷ Brain magnetic resonance imaging (MRI) has indicated diffuse white matter changes clearly associated not only with complete deficiency⁸⁻¹⁰ but also with partial deficiency of the laminin $\alpha 2$ chain.¹¹⁻¹³ This typical pattern is usually detectable after 6 months of age.¹⁴⁻¹⁵ Abnormal somatosensory and visual evoked potentials, as well as motor nerve conduction, have been described.¹⁶⁻¹⁷ Extensive brain abnormalities, such as cortical dysplasia, have been reported in some cases.⁴⁻¹⁸

Several mutations causing either a partial¹¹⁻¹⁹⁻²⁰ or a complete deficiency of the laminin $\alpha 2$ chain have been identified in the laminin $\alpha 2$ chain gene.²¹⁻²³ The laminins are a family of large trimeric basement membrane glycoproteins composed of α , β , and γ chains. The three subunit chains associate at their carboxyl termini in a coil, usually forming a cross shaped molecule with the long arm made up by the coil and the amino termini forming the short arms. The laminin-2 (merosin) and the laminin-4 (s-merosin) isoforms, which have molecular formulae $\alpha 2:\beta 1:\gamma 1$ and $\alpha 2:\beta 2:\gamma 1$, respectively, are characteristically enriched in basement membranes surrounding skeletal muscle fibres.²⁴ The high tissue specificity of these isoforms is provided only by the $\alpha 2$ chain, as the other component chains of the laminin isoforms, the $\beta 1$ or $\beta 2$ and $\gamma 1$ chains, are ubiquitous. The $\alpha 2$ polypeptide chain contains 3110 residues.²⁵ Detailed characterisation of the gene encoding this chain, LAMA2, showed that it exceeds 260 000 base pairs in size and contains 64 exons.²⁶

In the present study, we extended the sequencing of introns flanking all exons in order to enable PCR amplification of large enough segments for thorough mutation analysis of all the 64 exon regions. Oligonucleotide primers were designed to amplify each exon, which was then studied by PCR-SSCP analysis and direct sequencing. New mutations and several polymorphisms were identified and applications for prenatal diagnosis and search for founder effects were shown.

Department of Neuromuscular Diseases, Istituto Nazionale Neurologico "Carlo Besta", 20133 Milano, Italy
M Mora

Clinique de Pédiatrie et de Génétique Médicale, Groupe Hospitalier Pitié-Salpêtrière, 47 Boulevard de l'Hôpital, 75651 Paris Cedex 13, France
J-P Harpey

Hôpital des Enfants, Genève, Switzerland
C-A Haenggeli

Correspondence to:
Dr Guicheney.

Received 12 June 1997
Revised version accepted for publication 5 September 1997

Patients, material, and methods

PATIENTS

In the present study, mutation analysis or prenatal diagnosis was carried out on 14 families (nine consanguineous and five non-consanguineous) with at least one affected child with a clinicopathological diagnosis of CMD, according to the criteria of the International Consortium on Congenital Muscular Dystrophy.²⁷ These families originated from various countries: France (490, 1272, 1672, 2656, BE), Italy (4835, 4836, SI, LI), Turkey (1663, 1830), Israel (IF), Belgium (BU), and Morocco (2324). Some of them have been referred to us for prenatal diagnosis. All families gave informed consent before inclusion in this study. A muscle biopsy was performed on one affected child from each family and studied by immunohistochemistry, using an antibody against the C terminal part of the globular G domain of the human laminin $\alpha 2$ chain (Chemicon), as previously reported.⁵ Complete deficiency of the laminin $\alpha 2$ chain was observed in patients in 13 families, whereas partial deficiency was shown in one family (1830).

MICROSATELLITE TYPING

D6S407 and D6S1620 microsatellite markers were provided by Génethon human genetic linkage maps.²⁸ PCR amplifications were performed on 40 ng genomic DNA, as previously described.²⁹

SEQUENCING OF INTRONS AND PCR-SSCP ANALYSIS

Sequencing of all 64 exons and their immediate intronic boundaries (30 bp) has previously been reported.²⁶ Further sequencing of the introns was carried out directly on our previously purified λ clones or subcloned restriction fragments using exon specific oligonucleotide primers as well as the AmpliCycle kit (Perkin-Elmer). Oligonucleotide primers were designed to amplify each LAMA2 exon, based on the intronic sequences using the OLIGO 4.0 program. The primers were chosen so that at least 30 to 50 bp of flanking intron sequences were readable. Amplifications from genomic DNA were performed using the PCR "touchdown" method.³⁰ The primer pairs and the initial and final annealing temperatures are given in table 1, with decreasing of the annealing temperature by 1°C after each two cycles and with 30 cycles at lower temperature. At each cycle, denaturation was performed at 94°C for 40 seconds and extension at 72°C for one minute. Final extension was at 72°C for 2.5 minutes. For SSCP analysis, PCR products were denatured for two minutes at 95°C after dilution (1:4) in 10% saccharose buffer, according to Maruya *et al*³¹ with minor modifications, kept in ice, loaded onto 10% polyacrylamide (37.5:1) minigels, then run at 8 mA (per gel) at 7°C and 20°C in a Hoeffer apparatus. The bands were visualised after silver staining of the gels (Biorad). All abnormal conformers were sequenced. When polymorphisms were determined, allele frequencies

were estimated by the analysis of 200 normal chromosomes from unrelated subjects.

DIRECT SEQUENCING OF THE PCR PRODUCTS

PCR fragments were centrifuged through a Centricon 100 membrane (Amicon) and sequenced with a *Taq* cycle sequencing kit (Applied Biosystems) using fluorescent dideoxynucleotides and one of the PCR primers. Reaction products were run on an automated DNA sequencer (Applied Biosystems). All sequences were determined on both strands.

Results

DETERMINATION OF EXON FLANKING INTRON SEQUENCES AND DESIGN OF PRIMER PAIRS FOR MUTATION DETECTION

Subcloned restriction fragments of genomic λ phage clones containing the 64 exons were sequenced using cycle sequencing. At least 50 bp of intron sequences flanking each exon were determined to enable PCR amplification and sequencing of intron sequences likely to be essential for the splicing of the primary transcript. These sequences might be altered in some cases of CMD.

The intron sequences were used to design oligonucleotide primers (table 1) for PCR amplification of the exon regions of the LAMA2 gene. This gene contains exons with sizes varying between 6 and 270 bp.²⁶ Sixty-three primer pairs allowed a direct PCR amplification of an exon with the exon/intron boundaries. For exon 5, two overlapping primer pairs were designed. The sizes of the PCR products ranged between 129 and 345 bp, allowing SSCP analysis and direct sequencing.

IDENTIFICATION OF MUTATIONS IN THE LAMA2 GENE

Eight new mutations were identified in LAMA2 associated with MD CMD in nine families (table 2). M1 is a 2 bp deletion at position 1539 in exon 10, 1539delGT, in a French non-consanguineous family (1272). This deletion resulted in the formation of a stop codon (T[GT]AG→TAG) in the domain V that should produce a truncated protein of 496 residues. M2 is also a 2 bp deletion at position 2098 in exon 13, 2098delAG, resulting in a frameshift and a premature termination within 20 amino acid residues in domain IVb. This deletion was identified in three French non-consanguineous families (490, 1272, 1672) (figs 1 and 2). M3 is a C to A transversion at position 2950 in exon 20 changing a cysteine codon to a stop codon (TGC to TGA) in domain IIIb. This nonsense mutation, Cys967stop, was identified in two non-consanguineous families originating from Italy (LI, 4835) (figs 1 and 2). M4 is another nonsense mutation in exon 20 at position 3011, a C to T transition changing a glutamine codon to a stop codon (CAA to TAA). This mutation, Gln988stop, was identified in a consanguineous Italian family (4836). M5 is a 1 bp deletion at position 3171 in exon 21, 3171delG, resulting in a frameshift and a premature stop codon within 33 amino acid

residues in domain IIIb. This deletion was identified in a consanguineous Turkish family (1663). M6 is a C to A transversion at position 4687 in exon 31, changing a cysteine residue to a stop codon (TGC to TGA). This nonsense mutation in domain IIIa, Cys1546stop, was identified in a consanguineous Italian family (SI). M7 is a G to A transition at position 6916+1 of the consensus donor splice site of exon 47 in the DNA of the affected child of family 1672. It is the second mutation transmitted by the father. The mutation occurred in the carboxy terminal G globular domain and its exact effect at the protein level is not known. M8 is a 1 bp deletion at position 8314 in exon 58, 8314delA, in a French

consanguineous family (2656), resulting in a frameshift and a premature stop codon within five amino acids in the carboxy terminal G globular domain. The six first mutations should result in truncated proteins lacking domains I and II of the long arm of the laminin $\alpha 2$ chain which normally forms a coil with the carboxyl termini of the $\beta 1$ or $\beta 2$ and γ chains. The M7 and M8 mutations occurred in the carboxy terminal G globular domain. M8 should result in the truncation of the two last G repeat domains against which the commercial antibody is raised. The temperatures at which these mutations can be detected by SSCP, as well as the modifications of restriction site they induce, are given in table 2.

Table 1 Oligonucleotide primers and conditions for PCR amplification of LAMA2 exons

Exon	Position on cDNA	Exon size (bp)	Location upstream from exon	Forward primer sequence (5'-3')	Reverse primer sequence (5'-3')	Location downstream from exon	PCR product size	PCR conditions
1	50	112	-109	CCTCTTCCCCAGCAGCTG	CCAAGCCCCATGCCTCGAC	6	244	53°C
2	162	171	-25	CACCTTCATTTGTCCATC	GCCACACACCTACACACT	12	225	TD55-50°C
3	333	113	-24	CATATGATGCTGCTAACT	CCACAAACCACTAACTAA	60	214	TD45-40°C
4	446	243	-22	TGTTGTTGTTATACCTCCCT	TATCAAATGTGCTTAAATGG	19	306	TD55-50°C
5	689	270	-103	TGGGAGAATGGGAAGTTA	GCAAACATCATCAAGTCAG	Exonic	240	TD55-50°C
6	959	118	-59	GCTTCCTTTATGGTTCTA	ATGACGATATTTTGTGG	17	211	TD45-40°C
7	1077	179	-54	TAAATGTATCTGAAATCAAAT	GTTCAACTGTTTACAGAAGC	88	340	TD55-50°C
8	1256	100	-43	GTTTCTATTCACTTCGTT	AATCATATTGGGCTTTTA	45	205	TD44-39°C
9	1356	161	-40	TTGTTTTAGAAATGTTGA	TTACTGGAATAAACAAATG	4	222	TD44-39°C
10	1517	141	-56	CGCAGCTCATAATGTTGA	GCAGCAAACAGAACCAAA	15	229	TD50-45°C
11	1658	174	-57	AAAAGCTGCTGATAGATA	TCAGTTAAAGCAGAATAG	23	271	TD45-40°C
12	1832	102	-50	TGATTTAATAGCCCATCT	ACATCAGGAAGTAACACG	26	195	TD45-40°C
13	1934	212	-47	CTCTTTTCAGTTTACTC	CAAAGTAAATGTGTGAAC	27	303	TD44-39°C
14	2146	112	-34	ACTTATTCACCATCTCTT	AAGCACTAAGGATACATT	14	177	TD50-45°C
15	2258	114	-33	CCAAGACTGACTAAAGCC	CCAGGGTAGTAGCTGAAT	65	229	TD50-45°C
16	2372	128	-49	TGATCCCTGACACCAAAA	GAGCTTATTCATGGGGTA	15	209	TD50-45°C
17	2500	87	-54	CTTTGTGTATGTCCTTCT	AATCATTTTGCAAGACAG	17	175	TD45-40°C
18	2587	212	-65	AATATCTAAACATTGCC	ACATCAATTCTGTCAGGG	23	317	TD45-40°C
19	2799	107	-27	TGATCACAGGTCTCTCTT	TTATGGGTACAGCCTCTAC	8	159	TD50-45°C
20	2906	181	-52	ACTTCGAGTTAACTGATT	ATGTAGGACTTAAGTCAT	12	262	TD44-39°C
21	3087	137	-45	TTTCTATTTTCCCTTCTTTG	TATCAATGAGCAAAAAAGTGGT	9	212	TD55-50°C
22	3224	237	-38	AAAGTTGTTAATGGTTGC	GAAACAGAAATTGAGGGAG	33	325	TD44-39°C
23	3461	144	-79	CTCCCGTTATGCATTCTC	GGCTCAGCAGTTCCCTAC	5	245	TD50-45°C
24	3605	180	-77	CATGCAGTTCGTAACCTA	GATAAATCTCCAAATGGT	58	332	TD45-40°C
25	3785	189	-51	ACCACCTTGGAGAGCTTATC	CACCAACAAATGACTAACTT	4	263	TD55-50°C
26	3974	134	-40	CATTCAAGTTTGTCTATAG	AAAATAAGAGCTTGAATA	19	210	TD44-39°C
27	4108	118	-27	ATCTAAGTACATTCTCGC	CAGGTTAGTAGGAGGTAG	6	168	TD50-45°C
28	4226	135	-85	TGTCTTTGCAGCCACTGA	TTCAAGGAAGGCTCAGAG	7	244	TD50-45°C
29	4361	125	-29	ACAAACTTCTTCTCCCTT	GTTTCTGATTGGGAAATA	23	194	TD50-45°C
30	4486	87	-23	CAATCCTTTTCTTTCTGA	TGTGACCCCTTACCATAT	46	173	TD45-40°C
31	4573	194	-55	TGTGGAGACATGACTTGC	CAGCAACGAGTATACGT	3	269	TD55-50°C
32	4767	143	-54	TCTGCCTGGGATGTTTAG	CTGCCCTGCTTGTGACTG	12	226	TD50-45°C
33	4910	99	-26	ATGTTTATGGGATGGAAT	TAGGAAGAAGGTGATTGT	81	223	TD55-50°C
34	5009	112	-25	GTGTTTCCCGAATTGGT	ATGATGATTAAATGTGAT	13	167	TD55-50°C
35	5121	163	-47	TGGCATGTTTGTTTACTA	CTGGAATCTCAGTTTGT	32	259	TD50-45°C
36	5284	211	-58	ACCCTAAGGCAGTGACAT	CCCCACCTTCTGAGATTA	26	312	TD44-39°C
37	5495	117	-24	TGCTTGTTCATAATGGT	TCTCTTTTGTGTTTACC	11	169	TD44-39°C
38	5612	164	-25	AAATGCCCTCTTCTCTAC	CCAAAACAAATGACATAC	1	207	TD55-50°C
39	5776	139	-32	CCCTGCATACTGTTTTGAAT	GACTTCCATTCCCAGCACAT	18	208	TD55-50°C
40	5915	103	-21	CATTTGTTTCTTGTCCAC	TCCAATAGTGTGATAGAA	60	201	TD55-50°C
41	6018	117	-52	GCTGAGGGATGATAAAAC	GTTGGTAGTGCCCTGAATG	6	192	TD55-50°C
42	6135	183	-54	GCCCACTAACTCCACCC	ACACAATGAAAAAGGAGA	8	262	TD55-50°C
43	6318	6	-31	TTTGTTTTGCTTCCATGTGA	TTCTTTTACGCATCTACC	143	197	TD55-50°C
44	6324	155	-32	ATATACATGCACACTAAT	CAGTTCTGATATGACGAT	14	218	TD45-40°C
45	6479	144	-51	CAAACTTTCTGAGAGATT	AGCAGCCTAATGAAAAGT	17	229	TD50-45°C
46	6623	134	-34	TGATATCTCTTGTTTTTG	AATCTTTATTAAGTTGGT	4	189	TD44-39°C
47	6757	160	-30	TCCCCTTCACTTCAACAC	ACAGGAGGAGGATGAACA	7	214	TD50-45°C
48	6917	125	-21	TAACGGTATTTCTTCTG	ACCCTGGGAGAGTTCTCA	12	175	TD50-45°C
49	7042	163	-52	TGACAGACCGAATAGATA	CACGGAACCTCTGCTATG	15	247	TD44-39°C
50	7205	145	-29	CCACTGGGGTATGTTTAC	GGCCTATATTGCATTATT	16	207	TD44-39°C
51	7350	139	-52	ATGTGGTTGATATTGCTC	AACTTAACTCCTTAGCTTT	20	228	TD44-39°C
52	7489	12	-21	TTCTCTTTCCCGTTATCT	TGGTGTGCTCTGCTTCTG	78	129	TD55-50°C
53	7501	121	-33	ATTGCTTTTGCTTTTCTAT	AGCAGCCACATAAGTAA	50	221	TD50-45°C
54	7622	177	-29	TGTCTACTCTTCTTTTCT	TGCATTACAGCTAGGTGTG	7	234	TD55-50°C
55	7799	149*	-60	CTAAAGCTAAGCCATAAA	GGAAGTTCACCTGAGTTA	31	256	TD45-40°C
56	7947	177*	-51	TGTTTGAATCAGATGTG	GTAGTAGTAATGAGGAGA	10	255	TD45-40°C
57	8125	169	-27	AAAATCTTATTTATTACA	TCCCTTCTGAAATGACTC	51	264	TD44-39°C
58	8294	113	-71	TTAGACAGCATCATTACC	TCTTCATTATCGGTTCT	28	229	TD53-48°C
59	8407	190	-70	GATACCGCTCTATTTTAG	GTAATCCCTTAGGGTACT	21	298	TD45-40°C
60	8597	156	-57	TCTGCATATGTGAAATTT	TCAATAAATGAATCAGCC	38	268	TD45-40°C
61	8753	154	-86	TACACACATAGAGCAACC	TGGATCTAGCAAGAAGTT	52	309	TD44-39°C
62	8907	131	-40	ATCCTCTAATCCAAATA	ATCTACACATCAACAATA	41	229	TD44-39°C
63	9038	223	-66	TGTGTGAACCATCATGAT	GAAATTGTTGCTGGGGTA	13	319	TD50-45°C
64	9261	269	-24	GCCCTCTTGCAATTGCCTT	TCTGGAGTGCAATTAGCA	23	334	TD55-50°C

G at position 7947 is the last nucleotide of exon 55 and not the first of exon 56 as recently published by Zhang *et al.*²⁶ This induces size changes for exons 55 and 56 (*).

Table 2 New LAMA2 mutations causing laminin $\alpha 2$ chain deficiency

	Mutations	Exons	Domains	Sequence variation	Size of the putative proteins	SSCP	Restriction site
M1	Cys497stop	10	V	1539delGT	496	7°C	New <i>SpeI</i> and <i>MaeI</i>
M2	2098delAG	13	IVb	2098delAG	702	7 and 20°C	No change
M3	Cys967stop	20	IIIb	2950 C→A	966	7 and 20°C	No change
M4	Gln988stop	20	IIIb	3011 C→T	987	7°C	Loss <i>NlaIV</i> and <i>BanI</i>
M5	3171delG	21	IIIb	3171delG	1073	20°C	No change
M6	Cys1546stop	31	IIIa	4687 C→A	1545	20°C	Loss <i>PvuI</i> and <i>BsaAI</i>
M7	6916+1 G→A	47	G	6916+1 G→A	Unknown	7°C	No change
M8	8314delA	58	G	8314delA	2759	7 and 20°C	New <i>TaqI</i>

IDENTIFICATION OF LAMA2 POLYMORPHISMS

We detected 13 polymorphisms in nine exons (table 3). Eight were silent polymorphisms and five induced amino acid changes in exons 12, 37, 55, and 56. Several polymorphisms induce a loss or creation of a restriction site and all can be detected by SSCP analysis at 7°C or 20°C, or at both temperatures, as mentioned in table 3. The frequencies of the less frequent alleles were estimated for most of the polymorphisms (table 3). The A→G polymorphism at position 5551 in exon 37 is the most informative with a frequency of 38% for allele G and 62% for allele A. Several rare variants were identified in 2 to 5% of the chromosomes which were analysed. In exon 55, the presence of at least three polymorphisms accounts for the various SSCP patterns observed, each corresponding to a different allelic combination. The respective frequencies of these polymorphisms have not been determined. We identified three additional polymorphisms in introns 4, 30, and 52 in the PCR products corresponding to exons 5, 30, and 52, respectively.

APPLICATION TO RESEARCH OF FOUNDER EFFECTS

SSCP analysis of exon 13 and exon 20 PCR products showed the same deletion, 2098delAG, in three French families and the same nonsense mutation, Cys967stop, in two families originating from Italy (fig 1). These five families were non-consanguineous, suggesting that two different mutations cause the disease. We identified the two mutations in

only two families, family 1272 (2098delAG and 1539delGT) and family 1672 (2098delAG and 6916+1 G→A). Analysis of the microsatellite markers D6S407 and D6S1620, which are the closest so far known to the LAMA2 locus, did not show whether the mutations 2098delAG and Cys967stop were independent mutational events or related ones (fig 2).

By the study of intragenic polymorphisms, we were able to show that the three French families shared the same haplotype associated with the 2098delAG mutation, suggesting that these families probably have a remote degree of consanguinity (fig 3). The G at position 2848, which is the less common allele, is associated with the 2098delAG deletion in these families.

In the Italian families a rare variant of exon 37, A at position 5579, was found associated with the Cys967stop mutation. The two families shared the same haplotype, which also suggests a founder effect (fig 2).

APPLICATION TO PRENATAL DIAGNOSIS

Six families were referred to us for prenatal diagnosis (fig 3). In addition to the two microsatellites bordering the LAMA2 locus, D6S407 and D6S1620, we analysed at least exons 12, 19, 37, and 42 in each family for polymorphism detection. As shown in fig 3, by haplotyping the microsatellites and polymorphisms, we could determine the two at risk haplotypes in each family and their presence in the DNA samples from trophoblastic biopsies (fig 3). The polymorphisms of exons 12, 19, and 42, 1905 G→A, 2848A→G, and 6286 G→A, were informative in two, two, and one of the six families, respectively, whereas the polymorphism of exon 37, 5551 A→G, was informative in all the six families. For the cases described here, all the fetuses have received only one or no disease haplotype, without any evidence of recombination. In two cases, study of the mutations already identified (family 1272, see above, and family 2324, 2418delC in exon 16²³) have strengthened the conclusions of the haplotype analysis.

Discussion

The laminin $\alpha 2$ chain is a large protein of the extracellular matrix. The gene, LAMA2, contains 64 exons, 62 with sizes varying between 87 and 270 bp and two unusually small ones, exon 43 and exon 52, with 6 and 12 base pairs, respectively.²⁶ Further sequencing of exon flanking intron regions allowed PCR amplification of sufficiently long intron sequences to ensure sequencing and mutation detection of




Figure 1 SSCP analysis of the three French families with a 2098delAG mutation (A) and the two Italian families with a Cys967stop mutation (B). Abnormal conformers were detected for the probands compared to the control (C). In each family, the proband and one of his parents was heterozygous for the mutation in agreement with the haplotypes shown in fig 2.




Figure 2 Search for founder effects with intragenic polymorphisms. The haplotypes associated with the mutation 2098delAG found in French families (490, 1272, 1672) are in white boxes; those corresponding to the mutation Cys967stop in two Italian families (LI, 4835) are shown in grey boxes.

the splice consensus sequences. Optimal PCR conditions were determined for each exon region of the LAMA2 gene. By PCR-SSCP and direct sequencing, we identified 14 different mutations, including the eight new ones of this report, in CMD patients with complete merosin deficiency.^{21 23} All these mutations were 1 or 2 bp deletions, nonsense, or splice mutations, leading to truncated proteins and absence of immunoreactivity with antibody directed against the C-terminal part of the protein. Three additional mutations have been reported by other groups, including a 5 bp deletion^{20 22} and a large deletion.²² Systematic screening, exon by exon, of such a large gene is expensive and time consuming. Nevertheless, such an approach cannot be avoided to detect missense mutations, such as the Cys996Arg,

causing CMD with partial merosin deficiency.¹¹ To detect mutations causing CMD with a complete deficiency, the protein truncation test (PTT) would be a more appropriate approach,^{32 33} especially since we have only identified 60 to 70% of the mutations by PCR-SSCP mutation screening. By analysing the various LAMA2 exons, we identified 16 polymorphisms, three in introns and 13 in exons. Most of them are silent polymorphisms, but five induce an amino acid change without any known physiological consequences. Four of these polymorphisms, 1905G→A, 2848 A→G, 5551 A→G, and 6286 G→A, in exons 12, 19, 37, and 42, respectively, have an allele frequency above 16% and can be useful as polymorphic markers, especially for prenatal diagnosis. The A to G polymorphism

Table 3 LAMA2 polymorphisms

Exon	Nucleotide	Amino acid	Frequency	Restriction site	SSCP conditions
10	1540 T→C	497 TGT(Cys)→TGC(Cys)	2%	New PstI	7°C, 20°C
12	1847 G→A	600 GGA(Gly)→AGA(Arg)	2%	Loss MnlI	7°C
12	1905 G→A	619 CGT(Arg)→CAT(His)	18%	Loss MaeII	7°C
19	2848 A→G	933 CAA(Gln)→CAG(Gln)	24%	No change	7°C, 20°C
33	5005 C→G	1663 ACC(Thr)→ACG(Thr)	3%	Loss EcoRI	7°C
37	5551 A→G	1834 GAA(Glu)→GAG(Glu)	38%	New MnlI	7°C
37	5579 C→A	1844 CGT(Arg)→AGT(Ser)	2%	No change	7°C
42	6286 G→A	2079 ACG(Thr)→ACA(Thr)	16%	Loss BsmI	20°C
54	7669 C→G	2540 TCC(Ser)→TCG(Ser)	17%	No change	7°C
55	7809 T→C	2587 GTA(Val)→GCA(Val)	ND	No change	20°C
55	7879 C→G	2610 GTC(Val)→GTG(Ala)	ND	New BclI	20°C
55	7894 G→A	2615 CCG(Pro)→CCA(Pro)	ND	Loss HinfI	20°C
56	7955 A→G	2636 ACA(Thr)→GCA(Ala)	6%	No change	7°C

ND=not determined.




Figure 3 Haplotypes of informative polymorphisms and two microsatellites: application to prenatal diagnosis.

at position 5551 is the most informative one and induces the creation of a new *MnII* restriction site. It was informative in our experience for the 11 prenatal diagnoses we performed (data not shown). These polymorphisms can be analysed to obtain more information in case of recombination between the two flanking microsatellite markers. Some of them do not induce the creation or the loss of a restriction site but they can all be easily detected by SSCP. The pattern associated with each polymorphism can be established using a reference panel of DNA samples which have been sequenced.

We present here two new mutations in the LAMA2 gene in non-consanguineous families. A 2 bp deletion at position 2098 in exon 13, 2098delAG, which results in the truncation of the laminin $\alpha 2$ chain in domain IVb, was found in three French families. Analysis of intragenic polymorphisms suggests that these three families are related. The nonsense mutation, Cys967stop, in exon 20, which results in the truncation of the laminin $\alpha 2$ chain in domain IIb, was identified in two families of Italian origin. These two families also probably have a remote degree of consanguinity since a rare polymorphism, adenine at position 1847, was found linked to the mutation. The two families originate from the south of Italy on the Adriatic coast. Analysis of the various polymorphisms of the gene could be used to identify founder effects in the populations. We previously reported that the rare polymorphism in exon 10, cytosine at position 1540, was linked to the Gln1241stop nonsense mutation identified in a family originating from Tunisia.²¹

This study provides valuable methods for determining the molecular defects in LAMA2

causing merosin deficient congenital muscular dystrophy.

This work was supported by the Institut National de la Santé et de la Recherche Médicale, the Association Française contre les Myopathies (AFM, France), and the French Research Ministry (Actions Concertées Coordonnées - Sciences du vivant) as well as the Swedish Medical Research Council. We thank Dr A Urtizberea, Dr J C Lambert, Dr C Richelme, Dr M Medici, Dr M Shobat, Dr N Philip, and Dr J Leroy for referring families, and Dr V Paquis and Dr F Leturcq for DNA extraction.

- Fardeau M. Congenital myopathies. In: Mastaglia FL, Lord Walton of Detchant, eds. *Skeletal muscle pathology*. Edinburgh: Churchill Livingstone, 1992:237-81.
- Banker BQ. The congenital muscular dystrophy. In: Engel AG, Franzini-Amstrong C, eds. *Myology*. Vol 2. New York: McGraw-Hill, 1994:1275-89.
- Dubowitz V. 41st ENMC International Workshop on Congenital Muscular Dystrophy, Naarden, The Netherlands. *Neuromusc Disord* 1996;6:295-306.
- Pini A, Merlini L, Tomé FMS, Chevally M, Gobbi G. Merosin-negative congenital muscular dystrophy, occipital epilepsy with periodic spasms and focal cortical dysplasia. Report of three Italian cases in two families. *Brain Dev* 1996;18:316-22.
- Tomé FMS, Evangelista T, Leclerc A, et al. Congenital muscular dystrophy with merosin deficiency. *C R Acad Sci Paris* 1994;317:351-7.
- Minetti C, Bado M, Morreale G, Pedemonte M, Cordone G. Disruption of muscle basal lamina in congenital muscular dystrophy with merosin deficiency. *Neurology* 1996;46:1354-8.
- Osari S, Kobayashi O, Yamashita Y, et al. Basement membrane abnormality in merosin-negative congenital muscular dystrophy. *Acta Neuropathol* 1996;91:332-6.
- Philpot J, Sewry C, Pennock J, Dubowitz V. Clinical phenotype in congenital muscular dystrophy: correlation with expression of merosin in skeletal muscle. *Neuromusc Disord* 1995;5:301-5.
- Fardeau M, Tomé FMS, Helbling-Leclerc A, et al. Dystrophie musculaire congénitale avec déficience en mérosine: analyse clinique, histopathologique, immunocytochimique et génétique. *Rev Neurol (Paris)* 1996;152:11-19.
- Vainzof M, Marie SKN, Reed UC, et al. Deficiency of merosin (laminin M or 2) in congenital muscular dystrophy associated with cerebral white matter alterations. *Neuropediatrics* 1995;26:293-7.
- Nissinen M, Helbling-Leclerc A, Zhang X, et al. Substitution of a conserved cysteine-996 in a cysteine-rich motif of the laminin $\alpha 2$ -chain in congenital muscular dystrophy with partial deficiency of the protein. *Am J Hum Genet* 1996;58:1177-84.
- Herrmann R, Straub V, Meyer K, Wagner M, Voit T. Congenital muscular dystrophy with laminin $\alpha 2$ deficiency: identification of a new intermediate phenotype and

- correlation of clinical findings to muscle immunohistochemistry. *Eur J Pediatr* 1996;155:968-76.
- 13 Mora M, Moroni I, Uziel G, *et al*. Mild clinical phenotype in a 12-year-old boy with a partial merosin deficiency and central and peripheral nervous system abnormalities. *Neuromusc Disord* 1996;6:377-81.
 - 14 Mercuri E, Pennock J, Goodwin F, *et al*. Sequential study of central and peripheral nervous system involvement in an infant with merosin-deficient congenital muscular dystrophy. *Neuromusc Disord* 1996;6:425-9.
 - 15 Van der Knapp MS, Smit LME, Barth PG, *et al*. Magnetic resonance imaging in classification of congenital muscular dystrophies with brain abnormalities. *Ann Neurol* 1997;42:50-9.
 - 16 Mercuri E, Muntoni F, Berardinelli A, *et al*. Somatosensory and visual evoked potentials in congenital muscular dystrophy: correlation with MRI changes and muscle merosin status. *Neuropediatrics* 1995;26:3-7.
 - 17 Shorer Z, Philpot J, Muntoni F, Sewry C, Dubowitz V. Demyelinating peripheral neuropathy in merosin-deficient congenital muscular dystrophy. *J Child Neurol* 1995;10:472-5.
 - 18 Sunada Y, Edgar TS, Lotz BP, Rust RS, Campbell KP. Merosin-negative congenital muscular dystrophy associated with extensive brain abnormalities. *Neurology* 1995;45:2084-9.
 - 19 Allamand V, Sunada Y, Salih MAM, *et al*. Mild congenital muscular dystrophy in two patients with an internally deleted laminin α 2-chain. *Hum Mol Genet* 1997;6:747-52.
 - 20 Hayashi YK, Ishihara T, Domen K, Hori H, Arahata K. A benign allelic form of laminin α 2 chain deficient muscular dystrophy. *Lancet* 1997;349:1147.
 - 21 Helbling-Leclerc A, Zhang X, Topaloglu H, *et al*. Mutations in the laminin α 2-chain gene (LAMA2) cause merosin-deficient congenital muscular dystrophy. *Nat Genet* 1995;11:216-18.
 - 22 Pegoraro E, Mancias P, Swerdlow SH, *et al*. Congenital muscular dystrophy with primary laminin α 2 (merosin) deficiency presenting as inflammatory myopathy. *Ann Neurol* 1996;40:782-91.
 - 23 Guicheney P, Vignier N, Helbling-Leclerc A, *et al*. Genetics of laminin α 2 chain (or merosin) deficient congenital muscular dystrophies: from identification of mutations to prenatal diagnosis. *Neuromusc Disord* 1997;7:187-90.
 - 24 Wever UM, Engvall E. Merosin/laminin-2 and muscular dystrophy. *Neuromusc Disord* 1996;6:409-18.
 - 25 Vuolteenaho R, Nissinen M, Sainio K, *et al*. Human laminin M chain (merosin): complete primary structure, chromosomal assignment, and expression of the M and A chain in human fetal tissues. *J Cell Biol* 1994;124:381-94.
 - 26 Zhang X, Vuolteenaho R, Tryggvason K. Structure of the human laminin α 2-chain gene (LAMA2), which is affected in congenital muscular dystrophy. *J Biol Chem* 1996;271:27664-9.
 - 27 Dubowitz V. 22nd ENMC sponsored workshop on congenital muscular dystrophy held in Baarn, The Netherlands, 14-16 May 1993. *Neuromusc Disord* 1994;4:75-81.
 - 28 Dib C, Fauré S, Fizames C, *et al*. A comprehensive genetic map of the human genome based on 5264 microsatellites. *Nature* 1996;380:152-4.
 - 29 Helbling-Leclerc A, Topaloglu H, Tomé FMS, *et al*. Readjusting the localization of merosin (laminin α 2-chain) deficient congenital muscular dystrophy locus on chromosome 6q2. *C R Acad Sci Paris* 1995;318:1245-52.
 - 30 Don RH, Cox PT, Wainwright BJ, Baker K, Mattick JS. "Touchdown" PCR to circumvent spurious priming during gene amplification. *Nucleic Acids Res* 1991;19:4008.
 - 31 Maruya E, Saji H, Yokoyama S. PCR-LIS-SSCP (low ionic strength single-stranded conformation polymorphism)-a simple method for high resolution allele typing of HLA-DRB1, -DQB1, and -DPB1. *Genome Meth* 1996;6:51-7.
 - 32 Roest PAM, Roberts RG, van der Tuijn AC, *et al*. Protein truncation test (PTT) to rapidly screen the DMD gene for translation terminating mutations. *Neuromusc Disord* 1993;3:391-4.
 - 33 Roest PAM, Roberts RG, Sugino S, van Ommen GJB, den Dunnen JT. Protein truncation test (PTT) for rapid detection of translation-terminating mutations. *Hum Mol Genet* 1993;2:1719-21.